



Age	Children ≤15 years
Allergies	Propofol or severe egg allergy*** Allergy to muscle relaxants***
Health Status	<p>Patients from high level residential care</p> <p>Patients with disability requiring 24/7 care</p> <p>Patients from another hospital and/or ambulance transfer</p> <p>Patients who have severe neutropenia and recurrent infection</p> <p>Patients with diagnosed malignant hyperthermia or blood relatives with known malignant hyperthermia</p> <p>Patients with moderate/severe aortic mitral stenosis</p> <p>AMI or CVA in the last three months</p> <p>Elective cardiac stent insertion in the last 6 weeks not related to AMI/unstable angina</p> <p>Internal cardiac defibrillator in situ</p> <p>Patients with severe COPD / Emphysema / Asthma / Oxygen dependent</p> <p>Patients with severe renal impairment/having dialysis</p> <p>Patients with communicable diseases</p> <p>Patients who are pregnant</p> <p>Patients with history of difficult intubation, limited mouth opening <2 finger breadths</p> <p>Patients with moderate to severe limitation neck/jaw movement</p> <p>Patients with long COVID symptoms</p>
Weight / BMI	<p>>130kg maximum</p> <p>BMI of >40 or greater for Gastroscopy***</p> <p>BMI of >45 or greater for Colonoscopy***</p> <p><i>Note: If having both procedures, BMI for Gastroscopy must be followed.</i></p>
Mobility	<p>Requiring full assistance</p> <p>Unable to walk or get up on procedure trolley</p> <p>Patients who cannot tolerate laying on left side**/**</p>
Cognition / Behaviour	<p>Patients with unstable mental health conditions</p> <p>Patients with known history of self-harm or harm to others, aggression, substance abuse, confusion, delirium</p> <p>Patients with severe dementia</p>
Falls Risk	<p>Patients who are frail</p> <p>Patients who are on wheelchair</p>
Infection	Patients with infection requiring isolation precautions i.e. communicable diseases

** Please discuss with endoscopist

***Please discuss with anaesthetist +/- pre-admission anaesthetic consultation

Before contacting the anaesthetist, please have all relevant clinical information available.